

The puzzling clustering and bimodality of long GRBs optical afterglow luminosities

M. Nardini¹, G. Ghisellini², G. Ghirlanda²

¹ *SISSA/ISAS, Via Beirut 2-4, 34014 Trieste, Italy*

² *Osservatorio Astronomico di Brera, via E. Bianchi 46, I23807 Merate, Italy*

The study of the rest frame properties of long Gamma-Ray bursts (GRBs) afterglows is a fundamental aspect for a better understanding of the nature of these powerful explosions. The launch of the *Swift* satellite (November 2004) marked a strong improvement of the observational capabilities of X-rays and optical afterglows. We studied the intrinsic optical afterglows of a sample of long GRBs finding an unexpected clustering and a hint of bimodality of the optical luminosity distribution (at 12h after the trigger). Through a Montecarlo simulation we proved that both the observed clustering and bimodality are not simply due to selection effects but should hide important information in the understanding of the nature of the afterglow emission. These findings can shed also light on the nature of the large fraction of optically dark GRBs.

1 Optical luminosity distribution

The study of the long GRBs optical luminosity light-curves rest frame properties in pre-*Swift* era showed that the luminosity distribution of most GRBs clusters around a typical value. The optical luminosity distribution is much narrower than the observed flux distribution and is also narrower than the distributions of the prompt gamma ray energetics (Nardini et al. ¹; see also Liang & Zhang ²). We also found a hint of bimodality in the optical luminosity distribution since 3 under luminous GRBs were at more than 4σ below the brighter events mean luminosity.

The launch of the *Swift* satellite combined with the availability of a net of fast pointing ground based optical telescopes allowed a fast and rich optical follow-up starting from a few dozen of seconds after the trigger for a large number of events. In more recent works we verified whether the clustering and the bimodality obtained in the pre-*Swift* optical luminosity distribution of long GRBs is confirmed by more abundant data of the *Swift*-era. We selected a sample of all the long GRBs with known redshift, good optical photometry coverage at some hours after the trigger and with a published estimate of the host galaxy dust absorption A_V^{host} . As of the end of March 2008 we found a sample of 33 GRBs fulfilling our selection criteria (golden sample) and 20 without the A_V^{host} estimate. In the left panel of Fig. 1 we plot the rest frame K-corrected and de-reddened monochromatic luminosity light-curves at the central frequency of the *R* band in the rest frame time. At very early times (accessible now with *Swift*, and poorly sampled in pre-*Swift* era) the luminosities span some orders of magnitudes, but at later times we confirm both the clustering and bimodality observed in Nardini et al. ¹. The right panel of Fig. 1 shows the luminosity distribution at the common rest frame time of 12 h after trigger. Note the presence of a gap between the under luminous and the brighter events families (for more details see Nardini et al. ³).




Figure 1: *Left panel:* Rest frame L_{ν_R} light-curves of the *Swift* GRBs in our sample superposed to the pre-*Swift* ones. *Right panel:* Optical luminosity distribution at 12 h after trigger (rest frame time).

2 The role of selection effects and dark bursts

In order to test whether the observed clustering and bimodality are real or just due to observational biases we applied the method developed in Nardini et al.⁴. We analysed all the telescopes limiting magnitudes of the optically dark GRBs observations and we obtained the probability distribution for a GRB to be observed in the R band at 12h after trigger with a given limiting magnitude (TSF). Through a Montecarlo simulation we generated a large number of optical afterglows assuming i) a redshift distribution following the star formation rate (Porciani & Madau⁵), ii) different luminosity functions and iii) A_V^{host} distributions. We then check if they would be observable using the limiting magnitudes described by the TSF. The luminosity distribution of the observable simulated events was then compared with the real one.

We found that no unimodal intrinsic luminosity function can reproduce the real observed distribution. This can be reproduced either assuming an intrinsically bimodal luminosity function or assuming a strong achromatic absorption (grey dust) for about half of the simulated afterglows. In our simulation most of the GRBs belonging to the bright family events are usually observable even at large redshifts, while most of the faint family bursts are below the telescope sensitivity. The ones we observe are then the tip of the iceberg of a large family of optically faint bursts. The optically dark GRBs can therefore belong to these unobservable low luminosity events.

References

1. M. Nardini *et al*, A&A **451**, 821 (2006)
2. E. Liang & B. Zhang, ApJ **638**, 67 (2006)
3. M. Nardini *et al*, MNRAS **386**, L87 (2008)
4. M. Nardini *et al*, MNRAS **383**, 1049 (2008)
5. C. Porciani & P. Madau, ApJ **548**, 522 (2001)